

Lifetime Risk of Developing or Dying From Cancer

The lifetime risk of developing or dying from cancer refers to the chance a person has, over the course of his or her lifetime (from birth to death), of being diagnosed with or dying from cancer. These risk estimates are one way to measure of how widespread cancer is in the United States.

The following tables list lifetime risks of developing and dying from certain cancers for men and women in the US. The information is from the National Cancer Institute's Surveillance Epidemiology and End Results (SEER) database, and is based on incidence and mortality data for the United States from 2014 through 2016, the most recent years for which data are available.

The risk is expressed both in terms of a percentage and as odds.

- For example, the risk that a man will develop cancer pf the pancreas during his lifetime is 1.66%. This means he has about 1 chance in 60 of developing pancreatic cancer (100/1.66 = 60).
- Put another way, 1 out of every 60 men in the United States will develop pancreatic cancer during his lifetime.

These numbers are **average risks** for the overall US population. Your risk may be higher or lower than these numbers, depending on your particular risk factors.

Males

Risk of developing Risk of dying from By continuing to browse this site, you are accepting our terms of use, including the use of cookies. To read and understand > View Cookies √ Accept Terms & Cookies our Terms of Use and our use of cookies BEFORE you use our site, please review: Terms of Use

Esophagus	0.80	125	0.76	132
Hodgkin lymphoma	0.24	417	0.04	2,500
Kidney and renal pelvis	2.16	46	0.60	167
Larynx (voice box)	0.53	189	0.19	526
Leukemia	1.86	54	0.96	104
Liver and bile duct	1.44	69	1.02	98
Lung and bronchus	6.70	15	5.49	18
Melanoma of the skin	2.77	36	0.39	256
Multiple myeloma	0.93	108	0.47	213
Non-Hodgkin lymphoma	2.43	4 1	0.81	123
Oral cavity and pharynx	1.66	60	0.42	238
Pancreas	1.66	60	1.39	72
Prostate	11.60	9	2.44	41

By continuing to browse this site, you are accepting our terms of use, including the use of cookies. To read and understand our Terms of Use and our use of cookies BEFORE you use our site, please review:

Terms of Use

> View Cookies

	%	1 in	%	1 in
All invasive sites	38.70	3	18.33	5
Bladder (includes in situ)	1.18	85	0.35	286
Brain and nervous system	0.55	182	0.42	238
Breast	12.83	8	2.57	39
Cervix	0.63	159	0.22	455
Colon and rectum	4.08	25	1.68	60
Esophagus	0.24	417	0.20	500
Hodgkin lymphoma	0.20	500	0.03	3,333

By continuing to browse this site, you are accepting our terms of use, including the use of cookies. To read and understand our Terms of Use and our use of cookies BEFORE you use our site, please review:

Terms of Use

> View Cookies

Lung and bronchus	6.05	17	4.50	22
Melanoma of the skin	1.79	56	0.19	526
Multiple myeloma	0.71	141	0.38	263
Non-Hodgkin lymphoma	1.93	52	0.64	156
Oral cavity and pharynx	0.71	141	0.19	526
Ovary	1.25	80	0.88	114
Pancreas	1.60	63	1.35	74
Stomach	0.66	152	0.30	333
Thyroid	1.93	52	0.07	1,429

By continuing to browse this site, you are accepting our terms of use, including the use of cookies. To read and understand our Terms of Use and our use of cookies BEFORE you use our site, please review:

Terms of Use

> View Cookies

References

Howlader N, Noone AM, Krapcho M, Miller D, Brest A, Yu M, Ruhl J, Tatalovich Z, Mariotto A, Lewis DR, Chen HS, Feuer EJ, Cronin KA (eds). SEER Cancer Statistics Review, 1975-2016, National Cancer Institute. Bethesda, MD, https://seer.cancer.gov/csr/1975_2016/, based on November 2018 SEER data submission, posted to the SEER web site, April 2019.

Last Medical Review: January 13, 2020 | Last Revised: January 13, 2020

American Cancer Society medical information is copyrighted material. For reprint requests, please see our <u>Content Usage Policy</u>.

By continuing to browse this site, you are accepting our terms of use, including the use of cookies. To read and understand our Terms of Use and our use of cookies BEFORE you use our site, please review:

Terms of Use

> View Cookies